

# Variation of morphological traits and quality indices of micropropagated *Melia volkensii* clones before field planting

Constantin Dushimimana <sup>1,2\*</sup>, Titus Magomere<sup>2</sup>, Jackson Mulatya<sup>3</sup>, Jan Van den Abeele<sup>4</sup>, Florence Olubayo<sup>2</sup>, Guy Smaghe<sup>1</sup> and Stefaan P.O. Werbrouck<sup>1</sup>

<sup>1</sup> Department of Plants and Crops, Faculty of Bioscience Engineering, Ghent University, Coupure Links 653 and Valentin Vaerwyckweg 1, B-9000 Ghent, Belgium; guy.smaghe@ugent.be (G.S.); stefaan.werbrouck@ugent.be (S.P.O.W.)

<sup>2</sup> Department of Plant Science and Crop Protection, University of Nairobi, P. O. Box 29053-00625 Kangemi, Nairobi, Kenya; magomere.titus@ku.ac.ke (T.M.); olubayo@uonbi.ac.ke (F.O.)

<sup>3</sup> Kenya Forestry Research Institute, P.O Box 20412-00200 Nairobi, Kenya; jmulatya@kefri.org (J.M.)

<sup>4</sup> Better Globe Forestry, P.O Box 823-00606 Nairobi, Kenya; jan@betterglobeforestry.com (J.V.D.A.)

\* Correspondence: Constantin.dushimimana@ugent.be, +254790484424

**Abstract:** The quality of acclimatised *in vitro* cultivated plants is essential to ensure good survival and growth after planting in field conditions. After two months of acclimatization, this study revealed a significant variation in survival rate, shoot characteristics, root traits, and biomass features between 13 clones of *Melia volkensii*. Also, a number of quality indices such as the Dickson Quality Index (DQI), shoot dry weight: root dry weight ratio (S:R), and sturdiness quotient (SQ) showed a large variation. The survival rate was genotype-dependent, with an average of 85%. Extreme genotypes were tall, had long internodes, a lot of leaves, and a large leaf area. At the other side of the spectrum, there were small clones with short internodes and a reduced number of leaves and leaf areas. The high S:R values (>2) of the evaluated clones indicate an imbalance between the shoot and root system, negatively influencing the quality of the *in vitro* produced *M. volkensii* plants. The SQ for all clones was below the threshold value of 6, implying a good expected survival rate. This study illustrates that some clones might systematically score better for SQ, S:R, plant volume, and DQI after acclimatization and that these factors could form the basis for selection.

**Keywords:** Acclimatization; diversity; *in vitro* plants; *Melia volkensii*; roots; morphological

## 1. Introduction

*Melia volkensii* is a multi-purpose tree recommended for arid and semi-arid forestry. It is native to Kenya, Tanzania, and Somalia [1] and is propagated mainly by harvesting seeds from selected trees with open pollination, resulting in heterogeneous progeny. The ripening of the fruit

from flowering to fruiting takes about 12 to 13 months [2]. Moreover, extracting the tender seeds from the hard fruits is costly [3] and requires challenging skills. Plant tissue culture has the potential to significantly increase the supply of planting material in terms of quality and quantity. Despite previous work on *M. volkensii* *in vitro* propagation [4,5,6], information on the morphological variation between micropropagated clones after acclimatization is still scarce.

In recent years, our research team has refined the rooting and acclimatization protocols for *M. volkensii* clones [7]. A number of genotypes were selected for their good *in vitro* propagation, and the plants were successfully acclimatized in the greenhouse. This savannah tree must grow not only fast and reach a good length and stem thickness but also needs well-developed taproots to withstand dry periods and severe storms. To prevent uprooting, the taproot must also be well oriented. This study aimed to evaluate a number of clones and compare their morphological differences in order to determine whether, after acclimatization, they meet the standards set by farmers for establishment in the field. To this end, a combination of different above and below ground morphological characteristics was recorded and used to determine correlations between them and to calculate tree quality indexes.

## 2. Materials and methods

### 2.1. Initiation and multiplication

In this study, thirteen *M. volkensii* clones were used and maintained *in vitro*. Three clones were initiated from axillary buds of phenotype-selected trees from a progeny field trial conducted by the Kenya Forestry Research Institute (KEFRI) at Tiva (clones 20/21, E34, and E7), Kenya. Shoots of these trees were cut off and surface sterilized by rinsing in ethanol 70 % and incubating in a 20 % JIK® commercial bleach (3.5 % m/v Sodium hypochlorite) containing 0.005 % detergent (Teepol) for 15 min. After rinsing three times in sterile distilled water, meristems were excised under a binocular and transferred to test tubes containing 20 ml Murashige and Skoog's (MS) medium (Murashige & Skoog, 1962) supplemented with 30 g.l<sup>-1</sup> sucrose, 5 µM of meta-Topolin riboside (mTR), and 2 g.l<sup>-1</sup> gelrite.

The remaining clones were derived from seeds collected from phenotypically selected, established open-pollinated trees. Mak1 and Mak2 were selected *in vitro* from seeds harvested from a selected tree in a farmers' orchard in Kibwezi, Kenya. Kas5 (Kasigau origin) was obtained from KEFRI. The remaining clones (19007, 19011, 19016, 19003, 19015, 19008, and K.F1) were obtained from seeds from Better Globe Forestry, Kenya. After removing the pulp, the nut was cracked open, and seeds were selected with an intact seed coat. They were surface sterilised in the same way as the shoots. After the seed coat was cut lengthwise for scarification, they were transferred to test tubes containing MS medium supplemented with 30 g.l<sup>-1</sup> sucrose and 2 g.l<sup>-1</sup> gelrite. The pH of the medium was adjusted to 5.8 before autoclaving at 121°C for 15 min. Two weeks after germination, each seedling was divided

into nodes to start micropropagation on the same basal medium supplemented with 5  $\mu$ M mTR. Each subculture lasted four weeks.

## 2.2. Rooting

Shoots of four weeks old were cut in explants of approximately 1.5 cm long with two internodes and leaves. They were rooted using a modified McCown woody plant medium [8] with half concentrations of salts ( $K_2SO_4$  and  $MgSO_4$ ) and supplemented with 3% sucrose, 2 g.l<sup>-1</sup> gelrite, 0.02 M Silver Thiosulfate (STS), and 2  $\mu$ M Indole-3-butyric acid (IBA) (pH 5.8 before autoclaving). Per jar of 350 ml, 70 ml autoclaved media was dispersed. The jar had a 0.45  $\mu$ m PTFE filter cup (model TC-GR350, Shanghai Zeshine Equipment Co. Ltd., China). Five explants were transferred into each jar, and cultures were incubated in a growth chamber at 22  $\pm$  1 °C with a photoperiod of 16/8 h light/dark. A completely randomized design was used.

## 2.3. Acclimatization and Experimental design

The rooted *in vitro* shoots were manually stripped of the remaining adherent medium and washed with tap water. Then the plantlets were treated with 0.2% Pearl® 50 SC (Osho Chemical Industries Ltd) and planted in 300 ml pots with peat moss (Kekkilä LSM 2w R8264). The pots were covered with clear plastic cups for 12 days (Figure 1. a). The pots were watered three times a week. Through irrigation, fertilization with Rosasol NPK 19.19.19+ TE (Rosasol-Even) was applied at two-week intervals. The experiments were carried out in a greenhouse located at Kabete field station. The area is situated on 1° 14' 52" S latitude and 36° 44' 32" E longitude, at an altitude of 1846 m above sea level. The average temperature of the greenhouse was 27 °C.

## 2.4. Data collection

Data were collected two months after transplanting. Parameters recorded were plant height (PH) in cm, number of leaves per plant (NL), internode length (INL) in mm, number of leaflets per leaf (NLL), stem diameter (SD) in mm, leaf area (LA) in cm<sup>2</sup>, chlorophyll (SPAD value) (CHL), shoot fresh weight (SFW) and dry weight (SDW) in grams. Stem diameter was measured at the first leaf using a digital vernier caliper. Mean leaf area was determined from the top five leaves of each plant through ImageJ (<https://imagej.nih.gov/ij/>). Chlorophyll content was determined using a SPAD meter (Konica Minolta SPAD-502Plus, Japan). Root traits (number of roots per plant, root diameter, root length (cm), collar diameter (CD) in mm, root fresh (RFW) and dry weight (RDW) in grams, and leaf area were collected from 10 plants per clone from each repetition except for the one with low survival rate. Shoot and roots were oven-dried at 70°C for 72 hours before the shoot and roots dry weight determination. The shoot to root dry mass ratio (S:R) is an essential measure of seedling survival. It relates the evaporating surface to the water-absorbing surface. A healthy plant has a ratio between 1:1 and 1:2 [9,10]. The sturdiness quotient (SQ) was calculated as the ratio of plant height (cm) to collar diameter (mm). The smaller the SQ

value, the more robust the plant and the higher the expected survival rate, especially in windy or dry places. An SQ higher than 6 is undesirable [9]. The Dickson's Quality Index (DQI) reflects the plants' ability to survive and grow in the field, with performance increasing as the index values rise. It was calculated according to the following formula [11]:

$$DQI = \frac{\text{Plant Dry weight (g)}}{\frac{\text{Height (cm)}}{\text{Collar diameter (mm)}} + \frac{\text{Shoot dry weight (g)}}{\text{Root dry weight (g)}}}$$

Plants with an index greater than or equal to 0.2 [12] are considered to be of good quality. The aerial plant volume (V) was calculated using the following formula:

$$V = \frac{1}{3} \times \pi \times \left( \frac{CD}{2} \right)^2 \times H$$

Where

V= Aerial plant volume;

CD= Collar diameter (mm);

H= Plant height (cm).

## 2.5. Statistical analysis

Each treatment consisted of at least 30 plants per clone, and this was done in a completely randomized design (CRD). The experiment was repeated three times so that a total of 90 plants was used per clone. ANOVA was performed using IBM® SPSS® statistics (version 28). A post-hoc Duncan's multiple range test ( $p < 0.05$ ) was performed to separate the means.

## 3. Results

### 3.1. Shoot morphological traits

The plantlets were successfully acclimatized (Figure 1. b-d) with a mean survival rate ranging between 34% to 99%. There was a highly significant ( $p < 0.001$ ) difference in survival rate between the clones. Clone 19016 had the highest survival rate, while clone 19003 had the lowest. There were highly significant ( $p < 0.001$ ) differences for all shoot traits among elite *M. volkensii* clones after two months of acclimatization (Table 1). Clone plant height varied from 15.81 cm for 19008 to 5.94 cm for 19003. The lowest average number of leaves per plant (7.46) was recorded in clone 19003, while the highest number of leaves per plant (11.68) was recorded in clone 19016. The number of leaflets per leaf ranged from five (Figure 1. e) to seven (Figure 1. f). Three clones, namely 20/21, E7, and K.F1, had the highest number of leaflets per leaf, namely seven. The lowest number of leaflets per plant of 5.30, 5.64 and 5.64 were recorded from clone E34, 19008 and 19003, respectively. Internode length ranged from 1.44 mm for clone 19003 to 2.96 mm clone 19008. Clone 19008 had significantly the highest mean internode length. Chlorophyll content (SPAD meter) varied from 23.91 for KAS5 to 30.07 for 19003. There was no significant difference between the chlorophyll content of clone 19003 and other eleven clones, namely K.F1, E7, MAK1, MAK2, 19016, 19015, 19008, E34, 19007, 20/21 and 19011. The thickest stems, 3.6 and 3.4 mm,

were recorded with MAK1 and 19007, respectively, while 20/21 and 9015 had the smallest stem diameters, measuring 2.3 and 2.4 mm. Leaf area ranged from 16.99 cm<sup>2</sup> for 19003 to 41.87 cm<sup>2</sup> for 19016. Seven clones, namely 19003, 20/21, E34, MAK2, KAS5, E7 and 19011, had significantly smaller leaf areas than the other clones. Shoot fresh weight ranged from 1.95 g to 5.67 g for all clones. The SFW of clones 19016 and MAK1 was similar. The lightest shoots were recorded in clones 19003, 20/21, E34, KAS5 and E7 with 1.95, 2.09, 2.55, 2.99 and 3.01 g, respectively. Clone coded 19016 and MAK1 had the best SDW (1.14 and 1.05 g), while the lowest SDW was recorded for 20/21, 19003 and E34 with mean values of 0.32, 0.38 and 0.43 g, respectively.

**Table 1.** Mean values of survival rate and shoot traits variation among 13 elite *Melia volkensii* clones after acclimatization.

Clones code	Survival rate	PH (cm)	NL	NLL	INL (mm)	CHL
19015	95.0 ± 0.64 <sup>a</sup>	10.5 ± 0.55 <sup>cde</sup>	9.8 ± 0.18 <sup>b</sup>	5.9 ± 0.12 <sup>ef</sup>	1.9 ± 0.09 <sup>d</sup>	28.4 ± 0.71 <sup>a</sup>
19011	93.3 ± 6.67 <sup>a</sup>	12.2 ± 0.60 <sup>bc</sup>	10.3 ± 0.29 <sup>bc</sup>	6.4 ± 0.13 <sup>cde</sup>	2.5 ± 0.15 <sup>abc</sup>	26.2 ± 0.73 <sup>ab</sup>
20/21	84.0 ± 8.74 <sup>a</sup>	9.1 ± 0.42 <sup>de</sup>	8.2 ± 0.18 <sup>f</sup>	7.2 ± 0.14 <sup>a</sup>	2.1 ± 0.10 <sup>cd</sup>	26.4 ± 2.25 <sup>ab</sup>
K.F1	95.9 ± 2.41 <sup>a</sup>	10.6 ± 0.67 <sup>cd</sup>	9.9 ± 0.23 <sup>bc</sup>	7.0 ± 0.17 <sup>ab</sup>	2.0 ± 0.19 <sup>bcd</sup>	30.0 ± 0.99 <sup>a</sup>
KAS5	98.0 ± 1.96 <sup>a</sup>	9.1 ± 0.55 <sup>de</sup>	9.6 ± 0.26 <sup>cd</sup>	6.48 ± 0.16 <sup>bcd</sup>	1.9 ± 0.13 <sup>de</sup>	23.9 ± 0.71 <sup>b</sup>
MAK1	97.6 ± 1.19 <sup>a</sup>	12.0 ± 0.55 <sup>bc</sup>	9.7 ± 0.20 <sup>bcd</sup>	6.8 ± 0.09 <sup>abc</sup>	2.2 ± 0.11 <sup>bcd</sup>	29.1 ± 0.67 <sup>a</sup>
19008	83.8 ± 10.11 <sup>a</sup>	15.8 ± 0.85 <sup>a</sup>	9.8 ± 0.29 <sup>bcd</sup>	5.6 ± 0.18 <sup>fg</sup>	3.0 ± 0.18 <sup>a</sup>	27.9 ± 0.96 <sup>a</sup>
E34	58.0 ± 20.44 <sup>b</sup>	8.4 ± 0.82 <sup>e</sup>	9.0 ± 0.32 <sup>de</sup>	5.3 ± 0.13 <sup>g</sup>	2.1 ± 0.24 <sup>bcd</sup>	27.0 ± 0.87 <sup>ab</sup>
MAK2	95.9 ± 3.23 <sup>a</sup>	12.4 ± 0.87 <sup>bc</sup>	9.8 ± 0.23 <sup>bcd</sup>	6.5 ± 0.14 <sup>bcd</sup>	2.6 ± 0.22 <sup>ab</sup>	28.9 ± 0.83 <sup>a</sup>
19003	34.0 ± 9.52 <sup>c</sup>	5.9 ± 0.69 <sup>f</sup>	7.5 ± 0.51 <sup>g</sup>	5.6 ± 0.24 <sup>fg</sup>	1.4 ± 0.19 <sup>e</sup>	30.1 ± 1.14 <sup>a</sup>
19007	85.9 ± 6.49 <sup>a</sup>	13.6 ± 0.89 <sup>b</sup>	10.5 ± 0.29 <sup>b</sup>	6.2 ± 0.16 <sup>de</sup>	2.3 ± 0.17 <sup>bcd</sup>	26.6 ± 0.67 <sup>ab</sup>
19016	99.0 ± 1.01 <sup>a</sup>	14.1 ± 0.53 <sup>ab</sup>	11.7 ± 0.16 <sup>a</sup>	6.5 ± 0.09 <sup>cd</sup>	2.6 ± 0.12 <sup>abc</sup>	28.6 ± 0.53 <sup>a</sup>
E7	91.7 ± 4.79 <sup>a</sup>	10.7 ± 0.66 <sup>cd</sup>	8.6 ± 0.25 <sup>ef</sup>	7.0 ± 0.18 <sup>a</sup>	2.3 ± 0.16 <sup>bcd</sup>	29.2 ± 0.77 <sup>a</sup>
Mean	85.4	11.36	9.72	6.48	2.24	27.84
LSD 0.05	23.1	2.42	0.90	0.54	0.57	4.04

**PH:** Plant Height (cm). **NL:** Number of leaves per plant. **NLL:** Number of leaflets per leaf. **INL:** internode length (mm). **CHL:** chlorophyll content (SPAD meter). **SD:** stem diameter in mm. **LA:** Leaf area (cm<sup>2</sup>). **SFW:** shoot fresh weight. **SDW:** Shoot dry weight. **LSD:** Least significant differences of means at 5%. Means followed by the same letters within a column are not significantly different at  $p \leq 0.05$  (Duncan's multiple range test at 5%). Data are presented as mean ± Standard Errors. N=90.


**Table 1.** (Continued).

Clones code	SD (mm)	LA (cm <sup>2</sup> )	SFW(g)	SDW(g)
19015	2.4 ± 0.05 <sup>gh</sup>	29.8 ± 2.73 <sup>bcd</sup>	3.3 ± 0.25 <sup>cde</sup>	0.6 ± 0.05 <sup>def</sup>
19011	2.9 ± 0.06 <sup>def</sup>	25.9 ± 2.56 <sup>bcd</sup>	3.8 ± 0.38 <sup>cd</sup>	0.6 ± 0.06 <sup>de</sup>
20/21	2.3 ± 0.04 <sup>h</sup>	19.3 ± 1.47 <sup>e</sup>	2.1 ± 0.19 <sup>f</sup>	0.3 ± 0.03 <sup>g</sup>
K.F1	3.3 ± 0.06 <sup>bc</sup>	32.9 ± 2.93 <sup>b</sup>	4.1 ± 0.36 <sup>bcd</sup>	0.7 ± 0.07 <sup>cd</sup>
KAS5	3.1 ± 0.08 <sup>cd</sup>	24.8 ± 2.53 <sup>bcd</sup>	3.0 ± 0.31 <sup>def</sup>	0.6 ± 0.06 <sup>def</sup>
MAK1	3.6 ± 0.08 <sup>a</sup>	31.2 ± 2.48 <sup>bc</sup>	5.0 ± 0.35 <sup>ab</sup>	1.0 ± 0.08 <sup>ab</sup>
19008	2.9 ± 0.09 <sup>def</sup>	29.0 ± 3.04 <sup>bcd</sup>	3.8 ± 0.37 <sup>cd</sup>	0.9 ± 0.08 <sup>bc</sup>
E34	2.6 ± 0.08 <sup>fg</sup>	20.7 ± 3.55 <sup>de</sup>	2.5 ± 0.40 <sup>ef</sup>	0.4 ± 0.07 <sup>efg</sup>
MAK2	3.3 ± 0.10 <sup>bc</sup>	23.0 ± 2.51 <sup>cde</sup>	3.3 ± 0.31 <sup>cde</sup>	0.6 ± 0.05 <sup>de</sup>
19003	2.8 ± 0.16 <sup>ef</sup>	17.0 ± 2.59 <sup>e</sup>	1.9 ± 0.32 <sup>f</sup>	0.4 ± 0.09 <sup>fg</sup>
19007	3.4 ± 0.10 <sup>ab</sup>	31.4 ± 3.69 <sup>bc</sup>	4.3 ± 0.51 <sup>bc</sup>	1.0 ± 0.13 <sup>ab</sup>
19016	3.3 ± 0.10 <sup>bc</sup>	41.9 ± 2.74 <sup>a</sup>	5.7 ± 0.30 <sup>a</sup>	1.1 ± 0.06 <sup>a</sup>

E7	$2.9 \pm 0.08^{de}$	$25.5 \pm 2.68^{bcde}$	$3.0 \pm 0.28^{def}$	$0.6 \pm 0.05^{de}$
Mean	3.01	27.74	3.62	0.71
LSD 0.05	0.30	9.87	1.22	0.25

189 **SD:** stem diameter in mm. **LA:** Leaf area (cm<sup>2</sup>). **SFW:** shoot fresh weight. **SDW:** Shoot dry weight. **LSD:**  
190 Least significant differences of means at 5%. Means followed by the same letters within a column are not  
191 significantly different at  $p \leq 0.05$  (Duncan's multiple range test at 5%). Data are presented as mean  $\pm$  Standard  
192 Errors. N=90 except for SFW and SDW, where N=30.





**Figure 1.** Acclimatization of *in vitro* propagated *M. volkensii* clones. (a) Covered plantlets with clear pots after transplanting. (b–d) Full acclimatized *M. volkensii* plantlets. (e–f) Two different leaf morphology of *M. volkensii*; (e) leaf with five leaflets and (f) leaf with nine leaflets. (Scale bar = 1 cm).

### 3.2. Root morphology

The mean values for the number of roots per plant, root length, root diameter and collar diameter are presented in Table 2. Significant ( $p < 0.001$ ) variations were observed in root traits among clones (Figure 2). The highest number of roots per plant were recorded from clone 19016, followed by 20/21, K.F1, 19015 and 19007. Eight clones, namely 19008, E34, E7, 19003, MAK3, MAK1, 19011 and KAS5, recorded the lowest number of roots per plant after two months under acclimatization. Clone 19011 had the longest roots (11.6 cm) and E34 the shortest. The largest root diameter and collar diameter were measured at MAK1 (7.7 mm). Clones such as 20/21, E34, 19003, 19015 developed the smallest root diameter with mean values of 1.9, 2.1, 2.5 and 2.5


mm, respectively. The smallest collar diameter (CD) was recorded as 4.2, 4.3, 4.3 and 4.9 mm, respectively, from clones 19015, 20/21, E34, and 19003. The roots of MAK1 were significantly heavier compared to other clones. Contrastingly clone 20/21 was recorded as possessing the lowest RFW with a mean value of 1.0 g. The RDW varied between 0.1 g to 0.5 g for all clones. The heaviest RDW was recorded for MAK1, while the lightest were recorded for 20/21, E34, 19003, and 19015.

**Table 2.** Number of roots per plant, root length, root diameter, collar diameter and root biomass of elite *Melia volkensii* clones seedling after two months under acclimatization.

Clones code	NR	RL (cm)	RD (mm)	CD (mm)	RFW(g)	RDW(g)
19015	$6.8 \pm 0.66^{abc}$	$10.8 \pm 0.48^{abc}$	$2.5 \pm 0.14^d$	$4.2 \pm 0.13^e$	$1.0 \pm 0.08^e$	$0.1 \pm 0.01^{de}$
19011	$4.1 \pm 0.42^e$	$11.6 \pm 0.50^a$	$4.8 \pm 0.31^b$	$6.3 \pm 0.31^b$	$1.9 \pm 0.16^{bc}$	$0.2 \pm 0.02^b$
20/21	$7.5 \pm 0.60^{ab}$	$9.4 \pm 0.32^{ef}$	$1.9 \pm 0.11^d$	$4.3 \pm 0.17^e$	$0.5 \pm 0.04^f$	$0.1 \pm 0.01^e$
K.F1	$7.2 \pm 0.72^{ab}$	$10.3 \pm 0.48^{bcd}$	$4.2 \pm 0.19^{bc}$	$6.4 \pm 0.22^b$	$1.7 \pm 0.11^{cd}$	$0.2 \pm 0.01^{bc}$
KAS5	$4.1 \pm 0.47^e$	$10.0 \pm 0.46^{bcde}$	$4.7 \pm 0.26^b$	$5.6 \pm 0.16^{bc}$	$1.5 \pm 0.15^d$	$0.2 \pm 0.01^{bc}$
MAK1	$4.7 \pm 0.55^{de}$	$11.1 \pm 0.37^{ab}$	$7.7 \pm 0.43^a$	$7.7 \pm 0.43^a$	$3.8 \pm 0.17^a$	$0.5 \pm 0.03^a$
19008	$5.9 \pm 0.74^{bcde}$	$8.6 \pm 0.20^g$	$3.7 \pm 0.19^c$	$5.1 \pm 0.17^{cd}$	$1.4 \pm 0.10^d$	$0.1 \pm 0.01^{cd}$
E34	$5.8 \pm 0.64^{bcde}$	$8.1 \pm 0.32^g$	$2.1 \pm 0.15^d$	$4.3 \pm 0.19^e$	$1.0 \pm 0.12^e$	$0.2 \pm 0.01^e$
MAK2	$5.0 \pm 0.55^{cde}$	$9.8 \pm 0.31^{cdef}$	$4.0 \pm 0.29^{bc}$	$6.1 \pm 0.33^b$	$1.6 \pm 0.09^{cd}$	$0.2 \pm 0.01^{bc}$
19003	$5.0 \pm 0.81^{cde}$	$9.0 \pm 0.70^{efg}$	$2.5 \pm 0.27^d$	$4.9 \pm 0.28^{de}$	$1.0 \pm 0.17^e$	$0.1 \pm 0.01^e$
19007	$6.6 \pm 0.75^{abcd}$	$9.8 \pm 0.34^{def}$	$4.2 \pm 0.23^{bc}$	$6.0 \pm 0.19^b$	$2.0 \pm 0.17^{bc}$	$0.2 \pm 0.02^b$
19016	$8.4 \pm 0.66^a$	$8.7 \pm 0.21^{efg}$	$4.4 \pm 0.27^{bc}$	$5.7 \pm 0.16^{bc}$	$2.2 \pm 0.09^b$	$0.2 \pm 0.01^b$
E7	$5.5 \pm 0.55^{bcde}$	$10.0 \pm 0.44^{bcde}$	$3.6 \pm 0.23^c$	$5.8 \pm 0.21^{bc}$	$1.5 \pm 0.11^d$	$0.2 \pm 0.01^{bcd}$
Mean	5.95	9.87	3.98	5.62	1.67	0.18
LSD 0.05	2.20	1.41	0.90	0.87	0.44	0.06

NR: Number of roots per plant. RL: Root length (cm). CD: Collar diameter (mm). RFW: Root fresh weight. RDW: Root dry weight. LSD: Least significant differences of means at 5%. Means followed by the same letters within a column are not significantly different at  $p \leq 0.05$  (Duncan's multiple range test at 5%). Data are presented as mean  $\pm$  Standard Errors. N=30.





**Figure 2.** Morphological diversity of the root system of *M. volkensii* clones after two months of acclimatization under greenhouse conditions (scale bar = 1 cm).

### 3.3. Quality Indexes

There were highly significant ( $p < 0.001$ ) differences for SQ, DQI, and S:R among *M. volkensii* clones (Table 3). Sturdiness quotient ranged from 1.33 for clone 19003 to 2.74 for 19008. All clones had an SQ below the threshold of 6, which is recommended for robust young trees. The highest S:R ratio values of 7.58 were observed for 19015, while the lowest values of 2.35 and 3.45 were recorded for MAK1 and 19011. Since almost all evaluated clones had an S:R higher than the recommended value ( $\leq 1$ ), even MAK1, it can be concluded

that all acclimatized plants developed a disproportionately light root system in the pots. DQI index varied significantly among clones. The DQI ranged between 0.05 for 20/21 to 0.48 for MAK1. Seven clones out of thirteen had the average recommended value of  $\geq 0.2$ . The plant volume significantly varied among clones. Giant plants were observed in clones MAK1 with a mean shoot volume of 171.3, followed by clone 19007 (120.2), 19016 (119.3), K.F1 (113.1), and 19011, respectively. The plants with smaller volumes were recorded from 20/21 (41.6), 19003 (41.9), 19015 (47.7), and E34 (42.8), respectively. Clones ranking using plant volume index was comparable with the Dickson quality index ranking.

**Table 3.** Variation in quality indexes of 13 selected *Melia volkensii* clones after acclimatization.

Clones code	DQI	SQ	S:R	V
19015	$0.07 \pm 0.006^{de}$	$2.50 \pm 0.124^{abc}$	$7.58 \pm 0.600^a$	$47.7 \pm 3.68^d$
19011	$0.18 \pm 0.016^b$	$1.97 \pm 0.168^{cd}$	$3.45 \pm 0.291^{ef}$	$108.8 \pm 8.36^b$
20/21	$0.05 \pm 0.006^e$	$2.14 \pm 0.156^{bcd}$	$6.15 \pm 0.365^b$	$41.6 \pm 3.34^d$
K.F1	$0.16 \pm 0.010^{bc}$	$1.90 \pm 0.188^{cde}$	$4.48 \pm 0.567^{cde}$	$113.1 \pm 9.68^b$
KAS5	$0.17 \pm 0.020^{bc}$	$1.72 \pm 0.168^{de}$	$3.82 \pm 0.622^e$	$73.9 \pm 6.59^c$
MAK1	$0.48 \pm 0.051^a$	$1.75 \pm 0.160^{de}$	$2.35 \pm 0.234^f$	$171.3 \pm 13.43^a$
19008	$0.12 \pm 0.009^{cd}$	$2.74 \pm 0.219^a$	$6.14 \pm 0.457^b$	$95.3 \pm 8.31^{bc}$
E34	$0.06 \pm 0.008^{de}$	$2.09 \pm 0.222^{bcd}$	$5.73 \pm 0.401^{bc}$	$42.8 \pm 5.91^d$
MAK2	$0.16 \pm 0.016^{bc}$	$1.95 \pm 1.218^{cd}$	$4.10 \pm 0.380^{de}$	$96.1 \pm 7.91^{bc}$
19003	$0.07 \pm 0.013^{de}$	$1.33 \pm 0.145^e$	$5.57 \pm 0.975^{bcd}$	$41.9 \pm 6.35^d$
19007	$0.17 \pm 0.014^{bc}$	$2.27 \pm 0.239^{abcd}$	$4.71 \pm 0.378^{bcde}$	$120.2 \pm 10.80^b$
19016	$0.17 \pm 0.011^{bc}$	$2.59 \pm 0.161^{ab}$	$5.76 \pm 0.394^{bc}$	$119.3 \pm 7.90^b$
E7	$0.14 \pm 0.011^{bc}$	$1.80 \pm 0.193^{de}$	$4.41 \pm 0.405^{cde}$	$76.3 \pm 4.70^c$
Mean	0.16	2.08	4.88	91.6
LSD 0.05	0.07	0.66	1.62	29.14

**SQ:** Sturdiness quotient (Plant height to collar diameter ratio) (Value less than 6 indicates good quality plantlets [9]. **S:R:** Shoot to root ratio (S:R  $\leq 1$  indicates Good quality plantlets [10]. **DQI:** Dickson quality index (DQI  $\geq 0.2$  denote Good quality plants). **V:** Plant volume. **LSD:** Least significant differences of means at 5%. Means followed by the same letters within a column are not significantly different at  $p \leq 0.05$  (Duncan's multiple range test at 5%). Data are presented as mean  $\pm$  Standard Errors. N=30.

### 3.4. Correlation between morphological characteristics and quality of in vitro raised *Melia volkensii* plants

A strong correlation was recorded between several shoot and root parameters and quality indexes (Table 4). Plant height was strongly correlated ( $r=0.7-1$ ) with internode length and shoot dry weight and moderately associated with leaf area, plant height to collar diameter ratio and shoot fresh weight. The number of leaves per plant significantly correlated with leaf area, shoot fresh and dry weight. Stem diameter was strongly correlated with root diameter, collar diameter, root fresh, and dry weight. Average leaf area showed a strong correlation with the number of leaves per plant, shoot fresh and dry weight. Root diameter strongly correlated with collar diameter, root fresh, and dry weight. It was clear that shoot and root's

fresh and dry weight were significantly correlated. As a total dry weight, collar diameter, and RDW are positive elements of the Dickson quality index; they were obviously highly associated with it. However, SDW, a negative component of DQI, is still positively correlated with it. DQI was moderately correlated ( $r=0.5-0.7$ ) with root length and shoot fresh weight. Plant volume was strongly associated with SD, RD, DQI, SFW, SDW, RFW, and RDW. This correlation means that the higher the estimated plant volume, the higher the DQI, and the higher the plant biomass accumulation. Sturdiness quotient was moderately associated with a number of roots and internode length.

271

**Table 4.** Correlation between morphological characteristics and quality of in vitro raised *Melia volkensii* plants.

P	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																</
---	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	----

272 **SR:** Survival rate. **PH:** Plant Height. **NL:** Number of leaves per plant. **NLL:** Number of leaflets per leaf. **INL:** Internode length. **CHL:** Chlorophyll (SPAD  
273 value). **SD:** stem diameter. **LA:** leaf area. **NR:** Number of roots per plant. **RL:** Root length. **RD:** root diameter. **CD:** collar diameter. **DQI:** quality index. **SQ:**  
274 Plant Height to collar diameter ratio (Sturdiness quotient). **RFW:** root fresh weight. **RDW:** Root dry weight. **SFW:** shoot fresh weight. **SDW:** Shoot dry weight.  
275 **S:R:** shoot to root ratio. **V:** Plant volume. The green color indicates a positive correlation, and the red is negative. The darker the colors, the stronger the  
276 correlation, and the lighter the colors, the weaker.

## 4. Discussion

The quality of micropropagated plants after the weaning phase is essential for predicting survival and growth after field planting. In the present study, we assessed the phenotypic variation of a number of selected clones. We used these data to determine the correlations between the observed morphological parameters and calculate quality indices.

### 4.1. Survival

At the end of acclimatization, an average of 83% of the plants survived. The survival rate was genotype-dependent, as significantly more losses were recorded for E34 (field selection) and 19003 (seedling). Survival rate during acclimatization can be a valuable criterion for clonal selection. Similarly, the variation in the survival rate of acclimatizing Persian Walnut genotypes [13]. For *Uniola paniculata* [14], this variation has been attributed to the genotype-dependent degree of aberrant anatomy of the micropropagated plants. Similar survival rates were previously reported in other Meliaceae families, including *M. azedarach* L [15], *Khaya senegalensis* [16], *Toona ciliatae* [17].

### 4.2. Morphological characteristics of the shoots

Significant clonal differences were observed for shoot traits such as plant height, the number of leaves per plant and leaflets, internode length, stem diameter, leaf area, and shoot biomass among clones. This variation among *M. volkensii* clones is attributed to their genetic makeup [18,19, 20] and could form an additional basis for further selection. Clone 19016 and 19008 recorded the tallest plants with long internode compared to other clones; This indicates that they are superior in growth rate. Contradictory results were reported by [21], who observed no significant difference for plant height between *Ficus carica* L varieties. Plant height indicates access to sunlight and competition [22,23]. Tall plants have more photosynthetic capacity, and they are meant to resist weed competition [24,25].

The greatest number of leaves per plant and maximum leaf area were recorded for 19016. No significant differences were recorded among six clones out of 13, with the number of leaves per plant ranging between 9.57 and 10.27. Four fig tree varieties had a comparable number of leaves during acclimatization [21]. A large number of leaves and leaf areas indicates a high photosynthetic activity. Moreover, leaf area can be used partially for forecasting plant growth [26]. The clones coded MAK1 and 19007 had a larger stem diameter than the others. This can indicate rapid growth and a high survival rate after the outplanting of these clones to semi-arid conditions. Significant differences were observed among clones in chlorophyll (SPAD meter). The chlorophyll of *in vitro* propagated *Prunus africana* was reported to increase over time, and the photosynthesis rate was similar to the mother plants [27]. The variation in chlorophyll could be attributed to growth rate and genotypic similarity among clones.

### 4.3. Root traits



*Melia volkensii* seedling produces a thick carrot-like taproot that is essential for anchoring and storing water and nutrients during the dry period [28]. But the roots of the tissue-cultured trees branched before thickening. The average number of thick roots ranged from 4.1 to 7.2, depending on the genotype. The development of multiple taproots could, on the one hand, help plants to survive in semi-arid conditions but, on the other hand, reduce their root depth. This should therefore be determined through multi-year field trials. In *Hevea brasiliensis*, taproots and lateral roots development were attributed to genotypic differences between the clones [29]. A well-established seedling root system facilitates the fast establishment of plants after planting [30]. Root diameter and root length were significantly higher than 1.9 mm and 8.1 cm, respectively, indicating that the clones are genetically different in developing root traits. A more extensive root system is usually correlated with greater resistance to uprooting under heavy wind conditions [31]. The bigger the diameter of the roots, the more considerable 'carrot' root development. A combination of root length and lateral roots was applicable during the selection of excellent Norway spruce plants [32]. Root growth differences have been reported in *Fraxinus pennsylvanica* clones [33]. Seedlings with long roots and multiple secondary roots suggested a good water uptake and nutrient use efficiency [34]. Root collar diameter significantly varied with clones; although the minimal collar diameter was slightly above 4 mm while the maximum was nearly 8 mm. Our results are analogous to those of [35], who described variations in root collar diameter between clones of white spruce. *In vitro* plants with a larger collar diameter are considered superior because a robust rooting system promises high survival and growth rates after transplantation [24]. Clones significantly differed for root biomass. The heavy root was recorded from MAK1, suggesting that the more mass, the more chances to survive and grow better after planting [24,36].

#### 4.4. Quality indexes

The biomass and quality index of micropropagated *M. volkensii* clones differed significantly. Clone MAK1, followed by 19016 and 19007 respectively, were the best clones according to their SQ, S:R, plant volume, and DQI. The SQ for all clones was below the threshold value of six. This implies a good expected survival rate in dry areas [9]. The results showed that only one of the 13 clones, namely 'MAK 1', had an S:R value less than or equal to two. This confirmed its uniqueness. It indicates that for MAK1, the shoot is more or less in equilibrium with the root [10]. The SQ and S:R indices were developed for seedlings, and there is little literature on these indices applied to acclimatised *in vitro* plants.

The plant SQ indicates the vigor and robustness of the nursery tree plants. The clones with the highest robustness had more shoot biomass than roots, suggesting that they may be less resistant to extreme conditions [37], especially in semi-arid environments. After hardening off, transferring *in vitro* plants to large and long pots can be considered, which may improve the plants' S:R and quality. The planting volume increased in parallel with the increase in the DQI. Seven clones had high DQI values ( $\geq 0.2$ ). This indicates good quality; the higher this

index is, the better the survival and growth after transplanting [11,36,38].

#### 4.5. Correlations

Our results suggest that the large, heavy plants with many leaves and leaflets survived the best during the first critical month. Their final number of roots or root length did not play a role. During acclimatization of *Rhus coriaria* L., plantlets with high survival were longer, but unlike our results, they had more and longer roots [39].

The number of roots negatively correlated with root length, root diameter, root fresh and dry weight, collar diameter, and quality index, indicating that clones with more branched roots have shorter and thinner roots. More fibrous roots mean lower quality plants. The negative quality effect of a highly branched main root is not uncommon, as a good *M. volkensii* seedling is expected to have one deep taproot thickened like a carrot [28]. Our results showed that plant height to collar diameter ratio, known as SQ, had a slight negative correlation with stem and root diameter, root biomass, root length, chlorophyll, number of leaflets. The plant SQ increased parallelly with PH, NL, INL, NR, and LA. Our findings agree with [40], who reported a significant positive correlation between seedling height and SQ of *Castanea sativa* Mill. Populations.

S:R was negatively correlated with almost all parameters except root length, SQ, and chlorophyll. The high S:R values of evaluated clones indicate an imbalance between the shoot and root system, which negatively influences the quality of the *in vitro* produced *M. volkensii* plants. A low shoot-to-root biomass ratio between 1:1 to 1:2 indicates a vigorous plant [9] and a balanced morphological tree that can withstand storms. It also implies that water absorbed by the roots has equilibrium with water lost through transpiration. It also indicates that the amount of water absorbed by the roots is in balance with the transpirational area [41,10]. Moreover, DQI and V quality indexes of *M. volkensii* plants increased progressively with stem and root freshness and dry weight, suggesting that large biomass may be an indication of high quality. This was similarly found for *Acacia* spp, *Gliricidia sepium*, *Leucaena* spp [42], and *Anthocephalus cadamba* [43] whose seedlings' DQI was also associated with biomass.

## 5. Conclusion

The current study showed a significant variation of the phenotypic characteristics and the quality of *in vitro* produced plantlets between 13 selected *M. volkensii* clones. We demonstrated that, based on indices such as SQ, S:R, V, and DQI, a 'best' clone could be selected (MAK1), followed by 19011, 19016, 19007, KAS5, K.F1, and MAK2, respectively. Correlation analysis revealed that a good quality clone could be recognized by the higher stem diameter, plant biomass, root diameter, and collar diameter with a low shoot to roots ratio of its derived plants. We assume that these clones will provide insights into mass propagation and genetic improvement in the future. Selection of *M. volkensii* clones based on their quality after hardening off can be a good criterion before mass production. This will help to minimize losses after transplanting and during the tree's many years of growth under field conditions.

**Author Contributions:** Conceptualization, S.P.O.W., G.S., T.M., J.V.d.A. and C.D.; methodology, C.D. and S.P.O.W.; investigation, C.D.; resources, S.P.O.W. and G.S.; writing, original draft preparation, C.D. and S.P.O.W.; writing—review and editing, G.S., S.P.O.W., J.M., T.M., F.O. and J.V.d.A.; supervision, S.P.O.W., J.M., T.M.; project administration, S.P.O.W. and T.M.; funding acquisition, F.O., G.S., S.P.O.W. and T.M. All authors have read and agreed to the published version of the manuscript.

**Acknowledgments:** We greatly acknowledge Kenya Forestry Research Institute (KEFRI) and Better Globe Forestry Ltd for their assistance.

**Funding:** This work was supported by Belgian Development Cooperation through the VLIR-UOS team project “Synergy for mass propagation of elite *Melia volkensii* clones and exploiting its derived biopesticides” Ref KE2018TEA465A103.

## References

1. Stewart, M.; Blomley, T. Use of *Melia volkensii* in a Semi-Arid Agroforestry System in Kenya. *Commonw. For. Rev.* 1994, 73 (2), 128–131.
2. Orwa, C.; Mutua, A.; Kindt, R.; Jamnadass, R.; Simons, A. *Agroforestry Database: A Tree Reference and Selection Guide Version 4.0: Melia volkensii*. World Agroforestry Centre, Kenya; 2009.
3. Luvanda, A.; Musyoki, J.; Takeda, Y. *Melia volkensii* Enterprises for Enhanced Livelihoods in Semi-Arid Areas in Kenya. In Proceedings of International Conference on ‘Project on Development of Drought Tolerant Trees for Adaptation to Climate Change in Drylands of Kenya’ held at KEFRI Muguga 13–16 February 2017. KEFRI, Kenya; Ochieng, D., Kamondo, B., Muturi, G., Eds.; 2018; pp 117–128.
4. Indieka, S. A.; Odee, D. W.; Muluvi, G. M.; Rao, K. N.; Machuka, J. Regeneration of *Melia volkensii* Gürke (Meliaceae) through Direct Somatic Embryogenesis. *New For.* 2007, 34, 73–81. <https://doi.org/DOI10.1007/s11056-007-9038-7>.
5. Mulanda, E. S.; Adero, M. O.; Amugune, N. O.; Akunda, E.; Kinyamario, J. I. High-Frequency Regeneration of the Drought-Tolerant Tree *Melia volkensii* Gurke Using Low-Cost Agrochemical Thidiazuron. *Biotechnol Res Int* 2012, 2012, 818472. <https://doi.org/10.1155/2012/818472>.
6. Mulanda, E. S.; Adero, M. O.; Wepukhulu, D. K.; Amugune, N. O.; Akunda, E.; Kinyamario, J. I. Thidiazuron-Induced Somatic Embryogenesis and Shoot Regeneration in Cotyledon Explants of *Melia volkensii* Gürke. *Propag. Ornam. Plants* 2014, 14 (1), 40–46.
7. Werbrouck, S.; Bhogar, N.; Magomere, T.; Omondi, S. In Vitro Biotechnology of *Melia volkensii*, a High Potential Forestry Tree from Eastern Africa. In proceedings of the Fourth International Union of Forest Research Organizations (IUFRO) Unit 2.09. 02 on Development and application of vegetative propagation technologies in plantation forestry to cope with a changing climate and environment; 2017; Vol. 4, pp 249–256.
8. Lloyd, G.; McCown. Commercially-Feasible Micropropagation of Mountain Laurel, *Kalmia latifolia*, by Use of Shoot-Tip Culture. In *Int. Plant Prop. Soc. Proc.* 30; 1980; pp 421–427.
9. Jaenicke, H. *Tree Nursery Practices Practical Guide*; World Agroforestry Centre, 1999.
10. Haase, D. L. Morphological and Physiological Evaluations of Seedling Quality. In *TD National Proceedings: Forest and Conservation Nursery Associations-2006*. Proceedings RMRS-P-50. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. p 3-8, 50.; Riley, L., Dumroese, R., Landis, Eds.; 2007.
11. Dickson, A.; Leaf, A. L.; Hosner, J. F. Quality Appraisal of White Spruce and White Pine Seedling Stock in Nurseries. *For. Chron.* 1960, 36 (1), 10–13. <https://doi.org/10.5558/tfc36010-1>.
12. Nyoka, B. I.; Kamanga, R.; Njoloma, J.; Jamnadass, R.; Mng’omba, S.; Muwanje, S. Quality of Tree Seedlings Produced in Nurseries in Malawi: An Assessment of Morphological Attributes. *For. Trees Livelihoods* 2018, 27 (2), 103–117. <https://doi.org/10.1080/14728028.2018.1443027>.
13. Yegizbayeva, T. K.; García-García, S.; Yausheva, T. V.; Kairova, M.; Apushev, A. K.; Oleichenko, S. N.; Licea-Moreno, R. J. Unraveling Factors Affecting Micropropagation of Four Persian walnut Varieties. *Agronomy* 2021, 11 (7). <https://doi.org/10.3390/agronomy11071417>.
14. Aracama, C. V.; Kane, M. E.; Wilson, S. B.; Philman, N. L. Comparative Growth, Morphology, and Anatomy of Easy- and Difficult-to-Acclimatize Sea Oats (*Uniola paniculata*) Genotypes during in Vitro Culture and Ex Vitro Acclimatization. *J. Am. Soc. Hortic. Sci.* 2008, 133 (6), 830–843. <https://doi.org/10.21273/jashs.133.6.830>.
15. Vila, S.; Scocchi, A.; Mroginski, L. Plant Regeneration from Shoot Apical Meristems of *Melia azedarach* L. (Meliaceae). *Acta Physiol. Plant.* 2002, 24 (2), 195–199. <https://doi.org/10.1007/s11738-002-0011-y>.

16. Hung, C. D.; Trueman, S. J. In Vitro Propagation of the African Mahogany *Khaya senegalensis*. *New For.* 2011, 42 (1), 117–130. <https://doi.org/10.1007/s11056-010-9241-9>.
17. Li, P.; Shang, Y.; Zhou, W.; Hu, X.; Mao, W.; Li, J.; Li, J.; Chen, X. Development of an Efficient Regeneration System for the Precious and Fast-Growing Timber Tree *Toona ciliata*. *Plant Biotechnol.* 2018, 35 (1), 51–58. <https://doi.org/10.5511/plantbiotechnology.18.0130a>.
18. Runo, M. S.; Muluvi, G. .; Odee, D. W. Analysis of Genetic Structure in *Melia volkensii* (Gurke) Populations Using Random Amplified Polymorphic DNA. *African J. Biotechnol.* 2004, 3 (8), 421–425. <https://doi.org/10.5897/AJB2004.000-2080>.
19. Hanaoka, S.; Muturi, G. M.; Watanabe, A. Isolation and Characterization of Microsatellite Markers in *Melia volkensii* Gurke. *Conserv. Genet. Resour.* 2012, 4 (2), 395–398. <https://doi.org/10.1007/s12686-011-9558-5>.
20. Ndufa, J. K.; Miyashita, H. Preliminary Results on *Melia volkensii* Clonal Variation in Growth and Wood Properties in the Drylands of Kenya. In Proceedings of International Conference on ‘Project on Development of Drought Tolerant Trees for Adaptation to Climate Change in Drylands of Kenya’ held at KEFRI Muguga 13 -16 February 2017. KEFRI, Kenya Published; Ochieng, D., Kamondo, B., Muturi, G. M., Eds.; 2018; pp 90–95.
21. Bayoudh, C.; Labidi, R.; Majdoub, A.; Mars, M. In Vitro Propagation of Caprifig and Female Fig Varieties (*Ficus carica* L.) from Shoot-Tips. *J. Agric. Sci. Technol.* 2015, 17 (6), 1597–1608.
22. Gavin, D. G.; Peart, D. R. Spatial Structure and Regeneration of *Tetramerista glabra* in Peat Swamp Rain Forest in Indonesian Borneo. *Plant Ecol.* 1997, 131, 223–231. <https://doi.org/10.1023/A:1009771519028>.
23. Rijkers, T.; Pons, T. L.; Bongers, F. The Effect of Tree Height and Light Availability on Photosynthetic Leaf Traits of Four Neotropical Species Differing in Shade Tolerance. *Funct. Ecol.* 2000, 14 (1), 77–86. <https://doi.org/10.1046/j.1365-2435.2000.00395.x>.
24. Haase, D. L. Understanding Forest Seedling Quality: Measurements and Interpretation. *Tree Plant. Notes* 2008, 52 (2), 24–30.
25. Grossnickle, S. C.; MacDonald, J. E. Why Seedlings Grow: Influence of Plant Attributes. *New For.* 2018, 49 (1), 1–34. <https://doi.org/10.1007/s11056-017-9606-4>.
26. Weraduwege, S. M.; Chen, J.; Anozie, F. C.; Morales, A.; Weise, S. E.; Sharkey, T. D. The Relationship between Leaf Area Growth and Biomass Accumulation in *Arabidopsis thaliana*. *Front. Plant Sci.* 2015, 6 (APR), 1–21. <https://doi.org/10.3389/fpls.2015.00167>.
27. Komakech, R.; Kim, Y. G.; Kim, W. J.; Omujal, F.; Yang, S.; Moon, B. C.; Okello, D.; Rahmat, E.; Kyeyune, G. N.; Matsabisa, M. G.; Kang, Y. A Micropropagation Protocol for the Endangered Medicinal Tree *Prunus africana* (Hook f.) Kalkman: Genetic Fidelity and Physiological Parameter Assessment. *Front. Plant Sci.* 2020, 11 (November). <https://doi.org/10.3389/fpls.2020.548003>.
28. Kenya Forest Service. Guidelines to Growing *Melia volkensii* in the Dryland Areas of Kenya.
29. Carron, M. P.; Roux, Y. Le; Tison, J.; Dea, B. G.; Caussanel, V.; Clair, J.; Keli, J. Compared Root System Architectures in Seedlings and in Vitro Plantlets of *Hevea brasiliensis*, in the Initial Years of Growth in the Field. *Plant Soil* 2000, 223, 73–85.
30. Davis, A. S.; Jacobs, D. F. Quantifying Root System Quality of Nursery Seedlings and Relationship to Outplanting Performance. *New For.* 2005, 30 (2–3), 295–311. <https://doi.org/10.1007/s11056-005-7480-y>.
31. Albrecht, U.; Bodaghi, S.; Meyering, B.; Bowman, K. D. Influence of Rootstock Propagation Method on Traits of Grafted Sweet Orange Trees. *HortScience* 2020, 55 (5), 729–737. <https://doi.org/10.21273/HORTSCI14928-20>.
32. Högberg, K. A.; Bozhkov, P. V.; Von Arnold, S. Early Selection Improves Clonal Performance and Reduces Intracolonial Variation of Norway Spruce Plants Propagated by Somatic Embryogenesis. *Tree Physiol.* 2003, 23 (3), 211–216. <https://doi.org/10.1093/treephys/23.3.211>.
33. Kim, M. S.; Klopfenstein, N. B.; Cregg, B. M. In Vitro and Ex Vitro Rooting of Micropropagated Shoots Using Three Green Ash (*Fraxinus pennsylvanica*) Clones. *New For.* 1998, 16 (1), 43–57. <https://doi.org/10.1023/A:1006564423011>.
34. Batista, R. O.; Eduardo, A.; Neto, F.; Fernanda, S.; Deccetti, C.; Viana, C. S. Root Morphology and Nutrient Uptake Kinetics by Australian cedar Clones. *Rev. Caatinga* 2016, 29, 153–162. <https://doi.org/http://dx.doi.org/10.1590/1983-21252016v29n118rc>.
35. Lamhamedi, M. S.; Chamberland, H.; Bernier, P. Y.; Tremblay, F. M. Clonal Variation in Morphology, Growth, Physiology, Anatomy and Ultrastructure of Container-Grown White Spruce Somatic Plants. *Tree Physiol.* 2000, 20, 869–880. <https://doi.org/https://doi.org/10.1093/treephys/20.13.869>.
36. Tsakalimi, M.; Ganatsas, P.; Jacobs, D. F. Prediction of Planted Seedling Survival of Five Mediterranean Species Based on Initial Seedling Morphology. *New For.* 2013, 44 (3), 327–339. <https://doi.org/10.1007/s11056-012-9339-3>.
37. Takoutsing, B.; Tchoundjeu, Z.; Degrande, A.; Asaah, E.; Gyau, A.; Nkeumoe, F.; Tsoheng, A. Assessing the Quality of Seedlings in Small-Scale Nurseries in the Highlands of Cameroon: The Use of Growth

- Characteristics and Quality Thresholds as Indicators. *Small-scale For.* 2014, 13 (1), 65–77. <https://doi.org/10.1007/s11842-013-9241-7>.
38. Dina, K. D.; Pande, T.; Vladimir, M.; Mare, T. Dynamic of Growth and Quality of Arizona Cypress (*Cupressus arizonica* Greene) Seedlings from Three Container Types. In *Proceedings: International conference Reforestation Challenges*. 03-06 June 2015, Belgrade, Serbia. *Reforesta*; 2015. <https://doi.org/10.5281/zenodo.31739>.
  39. Amiri, S.; Mohammadi, R. Establishment of an Efficient in Vitro Propagation Protocol for Sumac (*Rhus coriaria* L.) and Confirmation of the Genetic Homogeneity. *Sci. Rep.* 2021, 11 (1), 1–9. <https://doi.org/10.1038/s41598-020-80550-4>.
  40. Tumpa, K.; Vidaković, A.; Drvodelić, D.; Šango, M.; Idžojić, M.; Perković, I.; Poljak, I. The Effect of Seed Size on Germination and Seedling Growth in Sweet Chestnut (*Castanea sativa* Mill.). *Forests* 2021, 12 (7), 1–11. <https://doi.org/10.3390/f12070858>.
  41. Bernier, P. Y.; Lamhamedi, M. S.; Simpson, D. G. Shoot: Root Ratio Is of Limited Use in Evaluating the Quality of Container Conifer Stock. *Tree Plant. notes* 1995, 46 (3), 102–106.
  42. Bayala, J.; Dianda, M.; Wilson, J.; Ouédraogo, S. J.; Sanon, K. Predicting Field Performance of Five Irrigated Tree Species Using Seedling Quality Assessment in Burkina Faso, West Africa. *New For.* 2009, 38 (3), 309–322. <https://doi.org/10.1007/s11056-009-9149-4>.
  43. Budiman, B.; Sudrajat, D. J.; Lee, D. K.; Kim, Y. S. Effect of Initial Morphology on Field Performance in White Jabon Seedlings at Bogor, Indonesia. *Forest Sci. Technol.* 2015, 11 (4), 206–211. <https://doi.org/10.1080/21580103.2015.1007897>.